

ORIGINAL ARTICLE

Impact of Saharan dust episodes on preterm births in Guadeloupe (French West Indies)

Jean-Francois Viel,¹ Yoann Mallet,¹ Christina Raghoumandan,² Philippe Quénel,³ Philippe Kadhel,⁴ Florence Rouget,¹ Luc Multigner³

Objectives Large amounts of mineral dust are transported from their African sources in the Saharan-Sahel region to the Caribbean Sea, generating peak exposures to particulate matter $\leq 10 \mu\text{m}$ (PM_{10}). This study aimed to investigate the impact of Saharan dust episodes on preterm births in the Guadeloupe archipelago.

Methods The study population consisted of 909 pregnant women who were enrolled in the TIMOUN mother-child cohort between 2004 and 2007. Desert dust episodes were assessed from PM_{10} concentrations recorded at the unique background air quality monitoring station located in Pointe-à-Pitre. For each woman, the daily PM_{10} concentrations were averaged over the entire pregnancy, and the proportion of days with intense dust episodes ($\geq 55 \mu\text{g PM}_{10}/\text{m}^3$) during pregnancy was calculated. Weighted logistic regression models adjusting for known individual sociomedical risk factors were used to estimate ORs and 95% CIs for preterm birth.

Results During pregnancy, the mean PM_{10} concentrations ranged from 13.17 to 34.92 $\mu\text{g}/\text{m}^3$, whereas the proportion of intense dust events ranged from 0.00% to 19.41%. Increased adjusted ORs were found for both the mean PM_{10} concentrations and the proportion of intense dust events (OR 1.40, 95% CI 1.08 to 1.81, and OR 1.54, 95% CI 1.21 to 1.98 per SD change, respectively). Restriction to spontaneous preterm births produced similar ORs but with wider 95% CIs.

Conclusion Considering the personal and social burden of this adverse pregnancy outcome, this finding is of importance for both healthcare workers and policy makers to provide necessary preventive measures.

INTRODUCTION

Desert dust is a natural contributor to atmospheric particulate matter (PM) worldwide. Large amounts of mineral dust are transported from their African sources in the Saharan-Sahel region to the Caribbean Sea.¹ The archipelago of Guadeloupe (French West Indies) is periodically exposed to desert dust between April and October, generating peak exposures that can last several days and exceed the $\text{PM}_{\leq 10 \mu\text{m}}$ (PM_{10}) health-based standards (WHO: 20 $\mu\text{g}/\text{m}^3$, European Union: 40 $\mu\text{g}/\text{m}^3$, as an annual average).

The available epidemiological evidence of the impact of desert dust episodes on human health is inconsistent. In Europe, although some studies have linked sandy dust storms to a range of adverse health outcomes including respiratory and cardiovascular conditions and mortality, others have

Key messages

What is already known about this subject?

- ▶ Large amounts of mineral dust are transported from the Saharan-Sahel region to the Caribbean Sea, generating peak exposures to particulate matter $\leq 10 \mu\text{m}$ (PM_{10}).
- ▶ The influence of desert dust particulate pollution on pregnancy outcomes appears plausible.

What are the new findings?

- ▶ This research adds new evidence to the health effects of Saharan dust episodes by showing an association between maternal pregnancy exposure (as assessed by the mean PM_{10} concentration and the proportion of intense dust events) and preterm birth while adjusting for sociodemographic and medical risk factors.

How might this impact on policy or clinical practice in the foreseeable future?

- ▶ Considering the personal and social burden of this adverse pregnancy outcome, this finding is of importance for both healthcare workers and policy makers to provide necessary preventive measures (such as communicating personalised information to pregnant women about a desert dust cloud moving to the coastline).

found no harmful effect.² The same inconsistencies emerge from studies published more recently.^{3–6} In the Caribbean region, a handful of studies is available.⁷ On the island of Barbados, no relationship was found between desert dust concentrations and asthmatic attendances at the Queen Elizabeth Hospital for the period of 1996–1997,⁸ whereas on the island of Trinidad, Saharan dust clouds were associated with increased acute paediatric asthma admissions.⁹ On the island of Martinique, PM_{10} concentrations were associated with increased hospital admissions for cardiovascular or pulmonary conditions over the period of 2001–2006.¹⁰ A recent study conducted in the Guadeloupe archipelago found a significant association between the PM_{10} and $\text{PM}_{2.5-10}$ pollutants contained in the Saharan dust and visits to the emergency department for children with asthma in 2011.¹¹ It is worth noting that these studies used an ecological design with time-series analyses, associating a number of

¹Univ Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, Rennes, France
²Gwad'air, Air quality monitoring agency, Petit-Bourg, France
³Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, Rennes, France
⁴Univ Antilles, CHU Guadeloupe, Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, Pointe-à-Pitre, France

Correspondence to

Pr. Jean-Francois Viel, Public Health, INSERM n° 1085, Rennes 35043, France; jean-francois.viel@univ-rennes1.fr

Received 3 August 2018

Revised 6 February 2019

Accepted 10 February 2019



© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Viel J-F, Mallet Y, Raghoumandan C, *et al.* *Occup Environ Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/oemed-2018-105405

health events on a particular day with air pollution levels on the same or recent days.

Regarding maternal exposure to anthropogenic particulate pollution (originating from the burning of fossil fuels in vehicles, power plants or various industrial processes), evidence from available studies is indicative of an association with preterm birth.¹² Therefore, the influence of desert dust particulate pollution on pregnancy outcomes appears plausible and could be of public health importance because Sahara dust transport may greatly increase the ambient levels of PM (particularly in Southern Europe and in some Atlantic islands).² Surprisingly, to the best of our knowledge, only one study has reported on the influence of Saharan dust episodes on pregnancy (in Spain).¹³

The Guadeloupe archipelago represents a unique opportunity to assess the potential impact of Saharan dust episodes on preterm birth for two main reasons. First, because of the absence of heavy industries and the year-round presence of trade winds (dispersing gaseous and chemical pollutants), anthropogenic particle pollution is low in this area (approximately 20 µg/m³).¹⁴ As a result, desert dust natural pollution is the main factor affecting PM₁₀ concentrations. Second, a high rate of preterm birth (15.8%) is observed in Guadeloupe.¹⁵ This high preterm birth rate is not fully understood, but it could be partly due to the African origin of its population, which as such presents a refractory high rate, common to populations of African descent, irrespective of their current geographic location and independent of socioeconomic status and obstetrical care.¹⁶ However, other than African ancestry, there is room for other unknown or unstudied risk factors (including environmental exposures) to explain this high rate of preterm births.

Therefore, this study aimed to investigate the impact of Saharan dust episodes on preterm birth among a population of pregnant women in the Guadeloupe archipelago using prospectively collected individual data.

METHODS

Setting and study design

Guadeloupe (part of the French West Indies) is an archipelago situated in the Caribbean Sea. It covers an area of 1628 km² and has a population of 450 000 inhabitants. This study relies on the TIMOUN mother-child cohort fully described elsewhere.¹⁵ Briefly, between 2004 and 2007, 1068 women attending check-up visits at public hospitals or dispensaries during their third trimester of pregnancy and having resided in Guadeloupe for >3 years were enrolled in the cohort. At inception, the participants completed a standardised questionnaire during a face-to-face interview with midwives. The questionnaire covered sociodemographic characteristics, medical and obstetrical history and various lifestyle factors. After delivery, information was collected from midwives, paediatricians and hospital medical records about the medical history of the pregnancy, delivery, perinatal conditions and measurements and health status of the newborn at birth. Following Rouget *et al*,¹⁵ we excluded women not born in the Caribbean (n=110), cases involving multiple births (n=25), severe birth defects (n=8) and induced pregnancies after fertility treatment (n=15), resulting in a study sample of 911 women (one case involved both fertility treatment and multiple births).

Preterm births

Gestational age in weeks was estimated by the obstetricians in charge of follow-up. It was based on the first day of the last menstrual period and was confirmed or corrected by ultrasound.

Preterm birth was defined as a birth before 37 completed weeks of gestational age. We distinguished between spontaneous and medically induced preterm birth. The latter results from the induction of delivery due to maternal or fetal factors (either by medication or by caesarean section) before the onset of spontaneous labour. Elective-induced labour was not practised at the delivery centres involved in this study.

Desert dust exposure assessment

Exposure data recorded at the urban background station located in Pointe-à-Pitre (central part of Guadeloupe, latitude 16°14'55" N, longitude 61°32'36" W) were provided by Gwad'Air (the official air quality monitoring agency for Guadeloupe), from 2005 to 2008. No other air quality monitoring station was operating in the Guadeloupe archipelago during this period of time. The median bird fly distance between the place of residence and the air quality monitoring station siting was 13.03 km (minimum 0.11 km, maximum 38.89 km).

Desert dust episodes were assessed in two ways. First, the mean daily PM₁₀ concentrations (µg/m³) were considered. However, as we could not distinguish between natural (desert dust) and anthropogenic concentrations, we also used a dust episode index, designed by Gwad'Air for daily transmission to local authorities and assumed to be more specific. It mainly relies on ATMO subindexes, daily indicators of air quality widely used in France.¹⁷ These subindexes, ranging from 1 (very good) to

Table 1 Distribution of sociodemographic and lifestyle factors in the study population (n=909, TIMOUN cohort, Guadeloupe archipelago, French West Indies, 2005–2008)

	N (%)
Maternal place of birth	
Guadeloupe or Martinique	795 (87.4)
Other Caribbean islands	114 (12.6)
Place of inclusion	
University hospital	605 (66.6)
Local hospital	222 (24.4)
Local antenatal care clinic	82 (9.0)
Maternal age (years)	
<20	74 (8.1)
20–34	535 (58.9)
>34	300 (33.0)
Marital status	
Single	229 (25.2)
Living with partner	466 (51.4)
Single living with family	189 (20.8)
Missing	25 (2.6)
Education (years)	
<5	62 (6.8)
5–12	663 (73.0)
>12	184 (20.2)
Body mass index (kg/m ²)	
<18.5	55 (6.1)
18.5–25.0	455 (50.0)
>25	379 (41.7)
Missing	20 (2.2)
Mother's employment during pregnancy	
Yes	378 (41.6)
No	530 (58.3)
Missing	1 (0.01)

Table 2 Descriptive statistics of desert dust exposure estimates during pregnancy on the entire study population and according to the mode of onset of labour (TIMOUN cohort, Guadeloupe archipelago, French West Indies, 2005–2008)

	N	Mean±SD	Minimum	Maximum
PM₁₀ concentrations (µg/m³)				
Whole cohort	909	27.15±3.08	13.17	34.92
Term birth	767	27.09±2.96	13.17	31.36
Preterm birth	142	27.49±3.68	14.82	34.92
Proportion of intense desert dust episodes (%)				
Whole cohort	909	7.73±4.06	0.00	19.41
Term birth	767	7.61±3.91	0.00	14.74
Preterm birth	142	8.34±4.77	0.00	19.41

10 (very poor), are calculated for each of four pollutants (sulfur dioxide, nitrogen dioxide, ozone and PM₁₀) on the basis of its concentration compared with a specific reference scale. Drawing on the ATMO PM₁₀ subindex category, the contrasts observed with the other subindexes, and the usual local meteorological conditions, Gwad'Air empirically defines a dust episode index as following: absent (0–27 µg PM₁₀/m³), light (28–38 µg PM₁₀/m³), moderate (39–54 µg PM₁₀/m³) and intense (≥55 µg PM₁₀/m³). As a categorical exposure estimate, we used the presence of an intense episode. Because of routine maintenance or machine failure independent of any desert dust episodes, there were a few missing days of data (14, 12, 21 and 13 in 2005, 2006, 2007 and 2008, respectively).

For each woman, daily measures of PM₁₀ concentrations were averaged over the entire pregnancy, and the proportion of days with intense dust episodes during pregnancy was calculated. However, we had to account for the lack of exposure data in the entire year of 2004, which included two full pregnancies (and 234 partial pregnancies). Therefore, we were forced to exclude those two pregnancies from the study sample. For the 909 remaining women, the number of days with available exposure measurements was used as the denominator of the intense dust episode proportion.

Covariates

In-depth statistical analyses have previously resulted in a list of individual risk factors for preterm birth in the TIMOUN cohort.¹⁵ The corresponding risk factors (all referring to the




Figure 1 Cumulative distribution of the mean particulate matter ≤10 µm (PM₁₀) concentration (µg/m³) (A) and the proportion of intense dust episodes (B) during pregnancy according to term (thick points)/preterm (thin points) births (TIMOUN cohort, Guadeloupe archipelago, French West Indies, 2005–2007).

index pregnancy) were considered in this study as covariates and introduced into the statistical models: place of inclusion (ie, the healthcare facility where pregnant women were enrolled: university hospital, local hospital, local antenatal care clinic), maternal age (<20, 20–34, >34 years), marital status (single, living as a couple, single and living with family), years of education (<5, 5–12, >12), body mass index (<18.5, 18.5–25.0, >25.0 kg/m²), parity (0, 1, ≥2), prior preterm birth (yes/no), prior miscarriage (yes/no), prior induced abortions (yes/no), lupus (yes/no), history of asthma (yes/no), chronic hypertension (yes/no), gestational weight gain (<275, 275–675, >675 g/week), gestational hypertension (yes/no), gestational diabetes (yes/no), urinary tract infection (yes/no) and sex of the newborn (male/female).

Data analysis

Multivariate logistic regression models were used to produce ORs and their 95% CIs for associations between desert dust exposure and preterm birth while adjusting for all individual risk factors previously described. As exposure estimates, averaged PM₁₀ concentrations and the proportion of intense dust episodes during pregnancy were considered in turn. To compare the impact of both exposure estimates, we calculated the ORs per SD change (3.08 µg/m³ and 4.06% for the mean PM₁₀ concentration and the proportion of intense dust episodes, respectively).

To compensate for differences in missing daily exposure measurements, each case in the input data set was weighted by the proportion of gestation length with available exposure measurements. Therefore, the more complete the exposure assessment, the more important the observation. We repeated analyses considering spontaneous and medically induced preterm births separately.

Statistical analyses were performed using R software (R Foundation for Statistical Computing, Vienna, Austria, 2018).

RESULTS

Characteristics of the study population

The women's pregnancies in this study occurred during the time period from 30 March 2004 to 30 July 2007. The sociodemographic and lifestyle factors of the study population are reported in table 1. Most women were enrolled in the study at the University Hospital, were aged between 20 and 34 years, were living with a partner and had completed between 5 and 12 years of education.

Approximately two-fifths were overweight or obese before pregnancy.

Regarding the outcome, 142 women (15.6%) delivered preterm among the 909 total women: 67 of them (47.2%) were spontaneous preterm births, and 74 (52.1%) were medically induced preterm births (for one woman, the type of preterm birth could not be identified).

Characteristics of exposure

The average PM₁₀ concentration observed during the time period of 2005–2008 was 26.78 µg/m³. Table 2 presents the distribution of desert dust exposure estimates during pregnancy for the entire cohort and according to the outcome (term/preterm birth). Mean PM₁₀ concentrations ranged from 13.17 to 34.92 µg/m³, whereas the proportion of intense dust events ranged from 0.00% to 19.41%. Figure 1 describes the cumulative distribution of both exposure estimates according to term/preterm births. Distributions are shifted to the right for preterm births, above 28 µg PM₁₀/m³ (figure 1A) and 7% of intense dust episodes (figure 1B).

Table 3 Crude and adjusted ORs for the risk of preterm birth according to desert dust exposure estimates and mode of onset of labour (TIMOUN cohort, Guadeloupe archipelago, French West Indies, 2005–2008)

Desert dust exposure estimates	No. of preterm births*	No. of term births	Crude		Adjusted†	
			OR‡	95% CI	OR‡	95% CI
All births	142	767				
PM ₁₀ concentrations (3.08 µg/m ³)			1.36	1.07 to 1.72	1.40	1.08 to 1.81
Proportion of intense desert dust episodes (4.06%)			1.38	1.11 to 1.72	1.54	1.21 to 1.98
Spontaneous preterm and term births	67	767				
PM ₁₀ concentrations (3.08 µg/m ³)			1.44	1.01 to 2.05	1.39	0.93 to 2.06
Proportion of intense desert dust episodes (4.06%)			1.53	1.11 to 2.10	1.63	1.13 to 2.35
Induced preterm and term births	74	767				
PM ₁₀ concentrations (3.08 µg/m ³)			1.28	0.93 to 1.76	1.30	0.91 to 1.86
Proportion of intense desert dust episodes (4.06%)			1.25	0.93 to 1.68	1.35	0.96 to 1.90

*For one woman, the type of preterm birth was unknown.

†The covariates for which we adjusted were place of inclusion, maternal age, marital status, years of education, body mass index, parity, prior preterm birth, prior miscarriage, prior induced abortions, lupus, asthma, chronic hypertension, gestational weight gain, gestational hypertension, gestational diabetes, urinary tract infection, sex of the newborn.

‡The ORs were calculated per SD change.

PM₁₀, particulate matter ≤10 µm; PTB, preterm birth.

Associations between preterm birth and desert dust exposure

Table 3 presents the crude and adjusted associations between desert exposure estimates and preterm births. Except for the association between spontaneous preterm births and mean PM₁₀ concentrations, the crude and adjusted ORs were very similar.

When considering all preterm births, increased adjusted ORs were found for both the mean PM₁₀ concentrations and the proportion of intense dust events (OR 1.40, 95% CI 1.08 to 1.81, $p=0.01$ and OR 1.54, 95% CI 1.21 to 1.98, $p=0.0006$, respectively). To determine whether these associations vary with distance to the monitoring station (potentially reflecting some exposure misclassification), we introduced an interaction term between distance (split at the median) and exposure estimates into the regression models. No significant interaction effects were observed ($p=0.75$ and $p=0.52$, for the mean PM₁₀ concentration and the proportion of intense dust episodes, respectively).

Restriction to spontaneous preterm births produced similar adjusted ORs but with wider 95% CIs (OR 1.39, 95% CI 0.93 to 2.06, $p=0.10$ and OR 1.63, 95% CI 1.13 to 2.35, $p=0.009$ for mean PM₁₀ concentrations and the proportion of intense dust events, respectively). Regarding induced preterm births, lower ORs were found, but they were still >1 (with CIs including unity).

DISCUSSION

This research adds new evidence to the health effects of Saharan dust episodes by showing an association between maternal pregnancy exposure (as assessed by the mean PM₁₀ concentration and the proportion of intense dust events) and preterm birth.

The present study has many strengths, including the study setting, the longitudinal design and extensive information on covariates. Guadeloupe can be considered a natural laboratory for disentangling the influence of desert dust natural pollution and anthropogenic pollution because according to Euphrasie-Clotilde *et al.*,¹⁴ on average, the concentration of anthropogenic pollution does not exceed 20 µg PM₁₀/m³ (despite some possible traffic-related pollution hot-spots). This study took advantage of the TIMOUN mother-child cohort, an archipelago-wide cohort, which collected information on potential maternal socioeconomic, behavioural and medical confounders at an early stage of pregnancy and therefore before outcomes occurred. To minimise residual confounding, we deliberately adjusted for numerous risk factors highlighted previously by Rouget *et al.*¹⁵

However, this study has some limitations. Exposure to Saharan dust was assessed through one single background air quality monitoring station (situated in the centre of Guadeloupe) regardless of the subject's residential location. These monitor-based estimates may have introduced some non-differential measurement error for pregnant women living far from the monitoring station due to the presumed spatial heterogeneity of air pollutants relating to topography and local micro sources, although the absence of interaction with distance to the monitoring station is reassuring. Alternative exposure estimates could be derived from desert dust modelling (such as the Navy Aerosol Analysis and Prediction System model) or from satellite imagery (such as the Moderate Resolution Imaging Spectroradiometer aerosol optical thickness product). However, considering the lower accuracy of these proxies and the need for retrospective exposure assessment (time period of 2004–2008), we preferred to rely on a well-established ground-based instrument. The population size and the number of events may appear modest but do not limit the statistical power of the analysis to identify significant findings.

In the Guadeloupe archipelago, the average PM₁₀ concentration observed during the time period of 2005–2008 (26.78 µg/m³) ranks high compared with those observed in the 17th largest French cities (mainland France) during the time period of 2007–2010 (range: 19.3–31.8 µg PM₁₀/m³).¹⁸ Associations are consistent across exposure estimates when calculating ORs per SD change suggesting that PM₁₀ concentrations recorded at the urban background monitoring station correctly reflect desert dust concentrations. Because the adjusted ORs do not contrast with the crude ORs, the potential for confounding is small. In other words, these results suggest that desert dust episodes influence preterm birth above and beyond the wide range of established risk factors. That higher ORs were found for spontaneous preterm births than medically induced preterm births is not unexpected. Although some underlying disorders are common to both outcomes (eg, preeclampsia), other specific conditions that cause induced preterm births are not assumed to be influenced by prenatal air pollution exposure, although a recent case-cross-over study suggested that a short-term increase in PM_{2.5} concentrations is a potential trigger of placental abruption.¹⁹

The underlying biological pathways of the effect of particulate pollution on pregnancy are complex and multiple, given

the chemical and physical heterogeneity of air pollutants and the various stages of fetal vulnerability during pregnancy. Researchers have, however, suggested the alteration of maternal-placental exchanges, endocrine disruption, oxidative pathways and alteration of maternal host-defence mechanisms as possible mechanisms.²⁰ However, compared with anthropogenic particle pollution, coarse particles coming from desert sources could bring their own toxicity because desert dust clouds carry heavy metals and biological components (such as bacteria, viruses, fungi and endotoxins) to the downwind countries.²¹

At variance with our results, Dadvand *et al*¹³ observed a small but statistically significant increase in gestational age at delivery in association with the number of episodic days during the third trimester and entire pregnancy. However, their study was conducted in another geographical region, with differences in climate pattern, distance from dust sources and morphology or chemical composition of the particles.

In conclusion, the current study suggests that exposure to desert dust episodes is associated with preterm birth, while adjusting for sociodemographic and medical risk factors. Considering the personal and social burden of this adverse pregnancy outcome, we believe that this finding is of importance for both healthcare workers and policy makers to provide necessary preventive measures (such as communicating personalised information to pregnant women about a desert dust cloud moving to the coastline).

Acknowledgements The authors would like to thank Sylvaine Cordier for implementing the TIMOUN cohort. The authors are grateful to Léah Michineau, Christine Monfort and Céline Garbin for valuable technical assistance. The authors would also like to thank their partners in the BrumiSaTerre research team (Stéphanie Guyomard, Antoine Talarmin, Séverine Ferdinand, Erika Upegui, Fabien Mercier and Gülen Ayhan) for fruitful discussions.

Contributors J-FV generated the idea for the paper and prepared an analytical plan. YM and J-FV conducted all data analyses. J-FV prepared the first draft on which all coauthors commented.

Funding This work was supported by the French National Research Program for Environmental and Occupational Health of ANSES (EST-2016/1/015).

Competing interests None declared.

Patient consent for publication Obtained.

Ethics approval This study was approved by the Guadeloupean Ethics Committee for studies involving human subjects.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data that support the findings of this study are available from the corresponding author on reasonable request.

REFERENCES

- Prospero JM, Mayol-Bracero OL. Understanding the transport and impact of African Dust on the Caribbean Basin. *Bull Am Meteorol Soc* 2013;94:1329–37.
- Karanasiou A, Moreno N, Moreno T, *et al*. Health effects from Sahara dust episodes in Europe: literature review and research gaps. *Environ Int* 2012;47:107–14.
- Neophytou AM, Yiallourous P, Coull BA, *et al*. Particulate matter concentrations during desert dust outbreaks and daily mortality in Nicosia, Cyprus. *J Expo Sci Environ Epidemiol* 2013;23:275–80.
- Alessandrini ER, Stafoggia M, Faustini A, *et al*. Saharan dust and the association between particulate matter and daily hospitalisations in Rome, Italy. *Occup Environ Med* 2013;70:432–4.
- Reyes M, Díaz J, Tobias A, *et al*. Impact of Saharan dust particles on hospital admissions in Madrid (Spain). *Int J Environ Health Res* 2014;24:63–72.
- Stafoggia M, Zauli-Sajani S, Pey J, *et al*. MED-PARTICLES Study Group. Desert dust outbreaks in Southern Europe: contribution to daily PM₁₀ Concentrations and Short-Term Associations with Mortality and Hospital Admissions. *Environ Health Perspect* 2016;124:413–9.
- Monteil MA. Saharan dust clouds and human health in the English-speaking Caribbean: what we know and don't know. *Environ Geochem Health* 2008;30:339–43.
- Prospero JM, Blades E, Naidu R, *et al*. Relationship between African dust carried in the Atlantic trade winds and surges in pediatric asthma attendances in the Caribbean. *Int J Biometeorol* 2008;52:823–32.
- Gyan K, Henry W, Lacaille S, *et al*. African dust clouds are associated with increased paediatric asthma accident and emergency admissions on the Caribbean island of Trinidad. *Int J Biometeorol* 2005;49:371–6.
- Blateau A, Bouobda D, Le Tertre A, *et al*. Health effects from desert dust episodes in Martinique, 2001–2006 [in French]. *Bull Veille Sanit Antilles Guyane* 2012;3:11–15.
- Cadelis G, Tourres R, Molinie J. Short-term effects of the particulate pollutants contained in Saharan dust on the visits of children to the emergency department due to asthmatic conditions in Guadeloupe (French Archipelago of the Caribbean). *PLoS One* 2014;9:e91136.
- Stieb DM, Chen L, Eshoul M, *et al*. Ambient air pollution, birth weight and preterm birth: a systematic review and meta-analysis. *Environ Res* 2012;117:100–11.
- Dadvand P, Basagaña X, Figueras F, *et al*. Saharan dust episodes and pregnancy. *J Environ Monit* 2011;13:3222–8.
- Euphrasie-Clotilde L, Molinie J, Feuillard T. *et al* The relationship between coastal West African dust level and Caribbean island dust. In: David AGD, Longhurst JWS, Brebbia CA, Barnes J, eds. *WIT transactions on ecology and the environment*. Southampton: WIT Press, 2017:121–8.
- Rouget F, Lebreton J, Kadhel P, *et al*. Medical and sociodemographic risk factors for preterm birth in a French Caribbean population of African descent. *Matern Child Health J* 2013;17:1103–11.
- Muglia LJ, Katz M. The enigma of spontaneous preterm birth. *N Engl J Med* 2010;362:529–35.
- van den Elshout S, Léger K, Nussio F. Comparing urban air quality in Europe in real time a review of existing air quality indices and the proposal of a common alternative. *Environ Int* 2008;34:720–6.
- Corso M, Pascal M, Wagner V, *et al*. Short-term impacts of particulate matter (PM₁₀) on mortality in 17 French cities, 2007–2010 [in French]. *Bull Epidémiol Hebd* 2015;1-2:14–20.
- Ananth CV, Kioumourtzoglou MA, Huang Y, *et al*. Exposures to air pollution and risk of acute-onset placental abruption: a case-crossover study. *Epidemiology* 2018;29:631–8.
- Slama R, Darrow L, Parker J, *et al*. Meeting report: atmospheric pollution and human reproduction. *Environ Health Perspect* 2008;116:791–8.
- Morman SA, Plumlee GS. The role of airborne mineral dusts in human disease. *Aeolian Res* 2013;9:203–12.